Contents

Volume 116 1992

ESSAY IN DEVELOPMENT		Otte, A. P., McGrew, L. L., Olate, J., Nathanson,	
Kimelman, D., Christian, J. L. and Moon, R. T. Synergistic principles of development: overlapping patterning system in <i>Xenopus</i>		N. M. and Moon, R. T. Expression and potential functions of G-protein a subunits in embryos of <i>Xenopus laevis</i>	141-146
mesoderm induction	1-9	Umbhauer, M., Riou, JF., Spring, J., Smith, J. C. and Boucat, JC.	
CONTRIBUTED PAPERS		Expression of tenascin mRNA in mesoderm during <i>Xenopus laevis</i> embryogenesis: the	
Kuhn, D. T., Sawyer, M., Packert, G., Turenchalk, G., Mack, J. A., Sprey, Th. E., Gustavson, E. and Kornberg, T. B.		potential role of mesoderm patterning in tenascin regionalization	147-157
Development of the <i>D. melanogaster</i> caudal segments involves suppression of the ventral regions of A8, A9 and A10	11-20	Narisawa, S. Hofmann, MC., Ziomek, C. A. and Millán, J. L. Embryonic alkaline phosphatase is expressed at M-phase in the spermatogenic lineage of the	
Smith, L. G., Greene, B., Veit, B. and Hake, S. A dominant mutation in the maize homeobox gene, <i>Knotted-1</i> , causes its ectopic expression in leaf cells with altered fates	21-30	mouse Kapur, R. P., Yost, C. and Palmiter, R. D. A transgenic model for studying development of	159-165
Rao, Y., Bodmer, R., Jan, L. Y. and Jan, Y. N.	21.00	the enteric nervous system in normal and aganglionic mice	167-175
The big brain gene of Drosophila functions to control the number of neuronal precursors in the peripheral nervous system	31-40	Goode, S., Wright, D. and Mahowald, A. P. The neurogenic locus <i>brainiac</i> cooperates with the <i>Drosophila</i> EGF receptor to establish the	
Brenner, H. R., Herczeg, A. and Slater, C. R. Synapse-specific expression of acetylcholine		ovarian follicle and to determine its dorsal- ventral polarity	177-192
receptor genes and their products at original synaptic sites in rat soleus muscle fibres regenerating in the absence of innervation	41-53	Collarini, E. J., Kuhn, R., Marshall, C. J., Monuki, E. S., Lemke, G. and Richardson, W. D.	
Wagner, M., Han, B. and Jessell, T. M. Regional differences in retinoid release from embryonic neural tissue detected by an in vitro		Down-regulation of the POU transcription factor SCIP is an early event in oligodendrocyte differentiation in vitro	193-200
reporter assay	55-66	Mullen, R. J., Buck, C. R. and Smith, A. M. NeuN, a neuronal specific nuclear protein in	
Ruiz i Altaba, A. Planar and vertical signals in the induction and patterning of the <i>Xenopus</i> nervous system	67-80	vertebrates Tilney, L. G., Cotanche, D. A. and Tilney, M. S.	201-211
Ruiz i Altaba, A. and Jessel, T. M. Pintallavis, a gene expressed in the organizer	07-00	Actin filaments, stereocilia and hair cells of the bird cochlea. VI. How the number and arrangement of stereocilia are determined	213-226
and midline cells of frog embryos: involvement in the development of the neural axis	81-93	Nieto, M. A., Bennett, M. F., Sargent, M. G. and	
Jägerbauer, EM., Fraser, A., Herbst, E. W., Kothary, R. and Fundele, R. Parthenogenetic stem cells in postnatal mouse		Wilkinson, D. G. Cloning and developmental expression of <i>Sna</i> , a murine homologue of the <i>Drosophila snail</i> gene	227-237
chimeras	95-102	Bunch, T. A. and Brower, D. L. Drosophila PS2 integrin mediates RGD-	000 047
French, V. and Brakefield, P. M. The development of eyespot patterns on butterfly wings: morphogen sources or sinks?	103-109	dependent cell-matrix interactions Krauss, S., Korzh, V., Fjose, A. and Johansen,	239-247
Schmidt-Ott, U. and Technqu, G. M. Expression of <i>en</i> and <i>wg</i> in the embryonic head		 Expression of four zebrafish wnt-related genes during embryogenesis 	249-259
and brain of <i>Drosophila</i> indicates a refolded band of seven segment remnants	111-125	Shi, DL., Feige, JJ., Riou, JF., DeSimone, D. W. and Boucat, JC.	
Hertzler, P. L. and Clark, W. H. Jr Cleavage and gastrulation in the shrimp Sicyonia ingentis: invagination is accompanied		Differential expression and regulation of two distinct fibroblast growth factor receptors during early development of the urodele amphibian	
by oriented cell division	127-140	Pleurodeles waltl	261-273

Contents

Delannet, M. and Duband, JL. Transforming growth factor-b control of cell- substratum adhesion during avian neural crest	275-287	Courtot, C., Frankhauser, C., Simanis, V. and Lehner, C. F. The <i>Drosophila</i> cdc25 homolog <i>twine</i> is required for meiosis	405-416
cell emigration in vitro	2/3-20/		403-410
Tabin, C. J.		Chandrasekhar, A., Ennis, H. L. and Soll, D. R. Biological and molecular correlates between induced dedifferentiation and spore germination	
Why we have (only) five fingers per hand: Hox genes and the evolution of paired limbs	289-296	in Dictyostelium	417-425
CONTRIBUTED PAPERS		Becker, T. and Macagno, E. R. CNS control of a critical period for peripheral induction of central neurons in the leech	427-434
Serbedzija, G. N., Bronner-Fraser, M. and Fraser, S. E.		Flamme, I. and Risau, W.	
Vital dye analysis of cranial neural crest cell migration in the mouse embryo	297-307	Induction of vasculogenesis and hematopoiesis in vitro	435-439
Yuan, J. and Horvitz, H. R. The Caenorhabditis elegans cell death gene ced-4 encodes a novel protein and is expressed during the period of extensive programmed cell		Goldstein, R. S. and Kalcheim, C. Determination of epithelial half-somites in skeletal morphogenesis	441-445
death	309-320	Tremml, G. and Bienz, M. Induction of <i>labial</i> expression in the <i>Drosophila</i>	
Andres, V., Nadal-Ginard, B. and Mahdavi, V. Clox, a mammalian homeobox gene related to Drosophila cut, encodes DNA-binding regulatory		endoderm: response elements for <i>dpp</i> signalling and for autoregulation	447-456
proteins differentially expressed during development	321-334	Waring, D. A., Wrischnik, L. and Kenyon, C. Cell signals allow the expression of a pre-	457 400
Freeman, M., Kimmel, B. E. and Rubin, G. M. Identifying targets of the <i>rough</i> homeobox gene of <i>Drosophila</i> : evidence that <i>rhomboid</i> functions in eye development	335-346	existent neural pattern in <i>C. elegans</i> Norbeck, B. A., Feng, Y. and Denburg, J. L. Molecular gradients along the proximal-distal axis of embryonic insect legs: possible guidance	457-466
Hainski, A. M. and Moody, S. A.		cues of pioneer axon growth	467-479
Xenopus maternal RNAs from a dorsal animal blastomere induce a secondary axis in host embryos	347-355	Cowing, D. W. and Kenyon, C. Expression of the homeotic gene <i>mab-5</i> during Caenorhabditis elegans embryogenesis	481-490
Conlon, R. and Rossant, J. Exogenous retinoic acid rapidly induces anterior ectopic expression of murine <i>Hox-2</i> genes in		Shamley, D. R., Opperman, L. A., Buffenstein, R. and Ross, F. P.	
vivo	357-368	Ontogeny of calbindin- D_{28k} and calbindin- D_{9k} in the mouse kidney, duodenum, cerebellum and	
Maeda, H., Yamagata, A., Nishikawa, S., Yoshinaga, K., Kobayashi, S., Nishi, K. and		placenta	491-496
Nishikawa, SI. Requirement of c-kit for development of intestinal pacemaker system	369-375	Geada, A. M. C., Gaunt, S. J., Azzawi, M., Shimeld, S. M., Pearce, J. and Sharpe, P. T. Sequence and embryonic expression of the murine <i>Hox-3.5</i> gene	497-506
Lüer, K. and Technau, G. M. Primary culture of single ectodermal precursors of <i>Drosophila</i> reveals a dorsoventral prepattern of intrinsic neurogenic and epidermogenic		Götz, M., Novak, N., Bastmeyer, M. and Bolz, J. Membrane-bound molecules in rat cerebral cortex regulate thalamic innervation	507-519
capabilities at the early gastrula stage DeFrances, M. C., Wolf, H. K., Michalopoulos,	377-385	Nishida, H. Regionality of egg cytoplasm that promotes	
G. K. and Zarneger, R. The presence of hepatocyte growth factor in the		muscle differentiation in embryo of the ascidian, Halocynthia roretzi	521-529
developing rat Leyser, H. M. O. and Furner, I. J.	387-395	Lallier, T., Leblanc, G., Artinger, K. B. and Bronner-Fraser, M. Cranial and trunk neural crest cells use different	
Characterisation of three shoot apical meristem mutants of <i>Arabidopsis thaliana</i>	397-403	mechanisms for attachment to extracellular matrices	531-541

687-695

gametogenesis

831-839

X inactivation in a subset of cells

Lamka, M. L., Boulet, A. M. and Sakonju, S. Ectopic expression of UBX and ABD-B proteins during <i>Drosophila</i> embryogenesis: competition, not a functional hierarchy, explains phenotypic suppression	841-854	Schulte-Merker, S., Ho, R. K., Herrmann, B. G. and Nüsslein-Volhard, C. The protein product of the zebrafish homologue of the mouse T gene is expressed in nuclei of the	
Doe, C. Q. Molecular markers for identified neuroblasts and ganglion mother cells in the <i>Drosophila</i> central nervous system	855-863	germ ring and the notochord of the early embryo Smith, D. E., Franco del Amo, F. and Gridley, T. Isolation of Sna, a mouse gene homologous to the Drosophila genes snail and escargot: its	1021-1032
Andres, A. J. and Cherbas, P. Tissue-specific ecdysone responses: regulation of the <i>Drosophila</i> genes <i>Eip28/29</i> and <i>Eip40</i>	005.050	expression pattern suggests multiple roles during postimplantation development	1033-1039
during larval development Artinger, K. B. and Bronner-Fraser, M. Notochord grafts do not suppress formation of neural crest cells or commissural neurons	865-876 877-886	Leder, A., Kuo, A., Shen, M. M. and Leder, P. In situ hybridization reveals co-expression of embryonic and adult α globin genes in the earliest murine erythrocyte progenitors	1041-1049
Shih, J. and Keller, R. The epithelium of the dorsal marginal zone of Xenopus has organizer properties	887-899	Friesel, R. and Brown, S. A. N. Spatially restricted expression of fibroblast growth factor receptor-2 during <i>Xenopus</i> development	1051-1058
Shih, J. and Keller, R. Cell motility driving mediolateral intercalation in explants of <i>Xenopus laevis</i>	901-914	González-Reyes, A., Macías, A. and Morata, G. Autocatalysis and phenotypic expression of Drosophila homeotic gene Deformed: its	
Shih, J. and Keller, R. Patterns of cell motility in the organizer and dorsal mesoderm of <i>Xenopus laevis</i>	915-930	dependence on polarity and homeotic gene function Martinez, S., Geijo, E., Sánchez-Vives, M. V.,	1059-1068
Weinmaster, G., Roberts, V. J. and Lemke, G. Notch2: a second mammalian Notch gene	931-941	Puelles, L. and Gallego, R. Reduced junctional permeability at interrhombomeric boundaries	1069-1076
Cui, X. and Doe, C. Q. <i>ming</i> is expressed in neuroblast sublineages and regulates gene expression in the <i>Drosophila</i> central nervous system	943-952	Dawe, R. K., and Freeling, M. The role of initial cells in maize anther morphogenesis	1077-1085
Bennett, R. L. and Hoffman, F. M. Increased levels of the <i>Drosophila</i> Abelson tyrosine kinase in nerves and muscles: subcellular localization and mutant phenotypes imply a role in cell-cell interactions	953-966	Griffith, C. M. and Hay, E. D. Epithelial-mesenchymal transformation during palatal fusion: carboxyfluorescein traces cells at light and electron microscopic levels	1087-1099
Hasan, G. and Rosbash, M. Drosophila homologs of two mammalian intracellular Ca ²⁺ -release channels: identification and expression patterns of the inositol 1,4,5-triphosphate and the ryanodine receptor genes	967-975	Donoghue, M. J., Patton, B. L., Sanes, J. R. and Merlie, J. P. An axial gradient of transgene methylation in murine skeletal muscle: genomic imprint of rostrocaudal position	1101-1112
Zimmer, A. and Zimmer, A. Induction of a RARβ2-lacZtransgene by retinoic acid reflects the neuromeric organization of the central nervous system	977-983	Bowerman, B., Tax, F. E., Thomas, J. H. and Priess, J. R. Cell interactions involved in development of the bilaterally symmetrical intestinal valve cells during	
Fischer-Vize, J. A., Rubin, G. M. and Lehmann, R. The fat facets gene is required for Drosophila eye and embryo development		embryogenesis in <i>Caenorhabditis elegans</i> Candia, A. F., Hu, J., Crosby, J., Lalley, P. A., Noden, D., Nadeau, J. H. and Wright, C. V. E.	1113-1122
Ekker, M., Wegner, J., Akimenko, M. A. and Westerfield, M. Coordinate embryonic expression of three zebrafish <i>engrailed</i> genes	1001-1010	Mox-1 and Mox-2 define a novel homeobox gene subfamily and are differentially expressed during early mesodermal patterning in mouse embryos Nieto, M. A., Gilardi-Hebenstreit, P., Charnay,	
Shimamura, K. and Takeichi, M. Local and transient expression of E-cadherin involved in mouse embryonic brain morphogenesis	1011-1019	P. and Wilkinson, D. G. A receptor protein tyrosine kinase implicated in the segmental patterning of the hindbrain and mesoderm	1137-1150

Weisshaar, B., Doll, T. and Matus, A. Reorganisation of the microtubular cytoskeleton by embryonic microtubule-associated protein 2 (MAP2c)	1151-1161	Legagneux, V., Bouvet, P., Omilli, F., Chevalier, S. and Osborne, H. B. Identification of RNA-binding proteins specific to Xenopus Eg maternal mRNAs: association with the portion of Eg2 mRNA that promotes deadenylation in embryos Hartenstein, A. Y., Rugendorff, A., Tepass, U.	•
Gould, A. P. and White, R. A. H. Connectin, a target of homeotic gene control in Drosophila	1163-1174		1193-1202
Nurcombe, V., Fraser, N., Herlaar, E. and Heath, J. K. MK: a pluripotential embryonic stem-cell-derived neuroregulatory factor	1175-1183	and Hartenstein, V. The function of the neurogenic genes during epithelial development in the <i>Drosophila</i> embryo	1203-1219
Brookman, J. J., Toosy, A. T., Shashidhara, L. S. and White, R. A. H. The 412 retrotransposon and the development of		1992 Index of Authors and Titles	1221-1234
gonadal mesoderm in <i>Drosophila</i>	1185-1192	1992 Subject Index	1235-1279